

Bacterial endophytes of aloe vera and their potential applications

Chitranshi Swati^{1*}, Gupta Akanksha¹, Sarkar Nagaji¹, Khare Priyanka¹

¹ College of Life Sciences, Cancer Hospital and Research Institute, Gwalior, M.P., India

*Corresponding author: swatichitranshi17@gmail.com

© The Author(s)

2022.

Published by ARDA.

Abstract

The study was aimed to assess culturable bacterial endophytes from the medicinal plant Aloe vera, their antimicrobial spectra against pathogens, and the potential of bacterial endophytes in textile and paper dyeing. Culturable seventeen bacterial endophytes were isolated from the Aloe vera plant out of which 16 showed varied antimicrobial activity against both human pathogens i.e., bacteria and fungi *E. coli*, *S. pyogenes*, acne bacterial isolate (ABI), *A. niger*, and *F. oxysporum*. Simultaneously, the bacterial endophyte ENDB3 is producing extracellular green-brown color pigment under submerged (SmF) condition and the extracted pigment has shown promising results in textile and paper dyeing at lab scale without using mordant. All the bacterial endophytes showed resistance against standard antibiotics (penicillin G P(10 units), Oxacillin (1 mcg), Cephalathin (30 mcg), Clindamycin (2 mcg), Erythromycin (15 mcg), and Amoxycylav (30 mcg)) at the specific concentration used. Concludingly, bacterial endophyte ENDB3 is found capable to produce bioactive molecules with pharmaceutical and dyeing industries which may provide a new path in the pursuit of new biological sources of drug and natural dyeing candidates. Hence, we suggest further evaluation and characterization of their bioactive molecules for pharmacological and dyeing potential.

Keywords: Bacterial endophytes, Antibacterial activity, Extracellular pigment

1. Introduction

Aloe vera is an evergreen medicinal plant of *Xanthorrhoeaceae* family. It is native to the Saudi Arabian Peninsula, sub-Saharan Africa and Western Indian Ocean islands. It is found in different type of climatic condition including coastal areas, grassland, and desert [1]. The Aloe plant has therapeutic and curative values hence, become the source for many potential uses. The exudates of fleshy leaves are widely used in cosmetic (Hair conditioner, moisturizers etc.) and medicinal purpose [2] as wound healing, burns, anticancer activity [3], diabetes and antimicrobial activity [4, 5].

Pathogenic microorganisms developed resistance to many commercially available medicines [6] which opens the door to find new bioactive compounds from natural sources. Recently, there is an emerging focus and interest in study of endophytes of medicinal plants. They are the alternative sources for bio- bioactive compounds. Endophytes are the microbial species, colonizing endosymbionts in inter and intra cellular parts of plants [7, 8]. It is hypothesized that because of the host-microbe relationship endophytes may become the source of many pharmacological bioactive compounds [7]. Hence endophytes and their bioactive compounds are known to be potent sources of novo natural products with possible exploitation and exploration in pharmacy and other industries. *Aloe vera* is a medicinal plant which provides a range of endophytic bacteria this becomes a motivating factor to explore their bioactive compounds for different industrial and medicinal applications. On the other hand, acne is also one of the serious issues of concern with many socio-health impacts which has to be solved, till today no product is suggested to control acne effectively. The bioactive molecules of bacterial

endophyte of *Aloe vera* can be used as a source of natural drug and in other potent industrial applications. Bioactive compounds can be used as natural colorant like food colorant, additives [9], natural dyes [10], cosmetic products, [11] and, functional foods [12] which represent an opportunity for the uses of natural pigments in different industry. For the production of natural pigment, the possibility to exploit microorganisms has been recommended [13]. In the present work endophytic bacteria from *Aloe vera* were isolated, and their bioactive molecules were screened for antimicrobial assay and extracellular pigment was explored for its industrial importance. This attempt is novel as application of extracellular pigments of endophytic bacteria from *Aloe vera* has not been reported, although antimicrobial activities of endophytic fungi and actinomycetes have been reported. In present study we have focused to isolate the endophytic bacterial flora of *Aloe vera* and attempts were made to find out the antimicrobial activity of endophytes against human pathogens and to explore screening of textile and paper dyeing potential of pigment produced by ENDB3.

2. Materials and methods

The *Aloe vera* plant sample was collected from the medicinal garden of College of Life Sciences, Cancer Hospital and Research institute, Gwalior (M.P.), India (geographically located in 26.2183° N and 78.18° E). Samples were immediately processed for analysis. Collected samples were washed with double distilled water and the samples were left for 10-15 minutes then the surface sterilization of sample was conducted by using 3% H₂O₂ washing for 3 min. followed with three times washing with sterile distilled water, excess water was allowed to drain. Plant sample cut into small pieces of size 2-3 cm soon after the single piece of collected plant sample inoculated on semisolid nutrient agar plate, incubated at 35±2°C for 48 hrs. BTB (Bromo thymol blue) was used as pH indicator.

Pus sample from acne was collected with sterile swab then spread, inoculated on nutrient agar plate. The plate was incubated at 35±2 °C for 24 hr. The pure isolate was obtained by following the method described above.

The pure colonies were obtained by subsequent rounds of quadrant streaking method on agar plates (1.5% W/V agar) was followed to obtain pure colonies of bacterial endophytes, incubation was held at 35±2 °C. The pure colonies were used for further studies. Similar was followed to obtain pure culture of acne pathogen.

Morphological parameters of each isolate such as colony size, shape, polysaccharide production, elevation, consistency, growth and colour were observed and recorded. Shape was determined by followed simple and Gram staining methods.

The antimicrobial potential of bacterial endophytes was evaluated against pathogenic bacterial and fungal strains included *Escherichia coli* MTCC-1302, *Streptococcus pyogenes* MTCC-1925, *Aspergillus niger* MTCC-281, *Fusarium oxysporum* MTCC-284 and acne bacterial isolate (ABI) pathogen. Kirby–Bauer disc diffusion method described by Bauer [14] was followed to perform the test. The pathogens were pre-cultured in Mueller Hinton (overnight for bacteria), potato dextrose broth (5 days for fungi), 0.5 ml of each broth culture was spread over agar plates then the sterile discs were impregnated with pre cultured bacterial endophytes, placed on pathogen seeded plates. The incubation was held at 35 ± 2°C for 48 h (bacteria) and 25°± 2°C C for 5 days (fungi), the zone of inhibition was determined. Antibiotic disc of tetracycline and nystatin were used as positive control for bacteria and fungi respectively. The experiment was performed in triplicates.

Antibiotic sensitivity pattern of bacterial endophytes was studied against different antibiotics where hexa disc of antibiotics (HiMedia) included penicillin G P (10 units), Oxacillin (1 mcg), Cephalathin (30 mcg), Clindamycin (2 mcg), Erythromycin (15 mcg) and Amoxyclav (30 mcg) was used. The pre-incubated bacterial endophyte culture was spread plated on MH agar medium then the antibiotic disc was placed and incubated at 35±2°C for 48 hr. Sensitivity of each isolate was recorded in terms of zone of inhibition produced [15].

The endophytic bacterial isolate ENDB3 is capable to produce extracellular green-brown pigment diffused in the agar medium therefore the attempts were made to assess its pigment production capabilities and their potential exploration in dyeing industries. 1 ml of pre-incubated overnight culture of selected bacterial strain (ENDB 3) was inoculated in 50 ml nutrient broth in an erlenmeyer flask incubated at 35 ± 2°C for 72 hr under static and unoptimized conditions. Optical density was recorded at 550nm after interval of 12 hr. The produced dark green-brown colour pigment was extracted by filtration followed by centrifugation at 10,000 rpm, performed in triplicates.

For assessment of dyeing potential, the extracted pigment was applied to paper, cotton and cotton cloth pieces. The plates were left for 24hr under soaking condition without mordant after soaking the dyed objects were washed with water and allowed to air dry at room temperature.

Data were analyzed by one-way analysis of variance (ANOVA) to determine statistical significance. A p -value of ≤ 0.05 was considered to be statistically significant.

3. Results

The results showed that the *Aloe vera* plant enriched with bacterial endophytes, total 17 different isolates namely, ENDB1 to ENDB17 were isolated from *Aloe vera* plant sample and identified on the basis of macro and microscopic attributes (see Table 1). All the isolated culturable bacterial endophytes are gram positive in nature and rod in shape.

Table 1. Morphological characteristics of bacterial endophytes

Strain No-	Shape	Size	Elevation	Polysaccharide production	Colour	Consistency	Growth	Gram Staining
ENDB1	Rod	Pin point	Convex	No	Creamish	Translucent	Slow	positive
ENDB2	„	„	„	Low	Creamish	„	„	„
ENDB3	„	Small	„	No	Green-Brown	„	Fast	„
ENDB4	„	Pin point	Flat	Low	Yellowish-white	„	„	„
ENDB5	„	Small	Flat and rough	High	White	„	„	„
ENDB6	„	Small	Flat	Low	Creamish	„	„	„
ENDB7	„	Pin point	Flat	Low	White	„	„	„
ENDB8	„	Small	Convex	High	Creamish	„	„	„
ENDB9	„	Pin point	Flat	Low	Creamish	„	„	„
ENDB10	„	Pinpoint	Convex	Low	White	„	„	„
ENDB11	„	Moderate	Convex	High	Creamish	„	„	„
ENDB12	„	Pin point	Raised	High	Creamish	„	„	„
ENDB13	„	Pin point	Convex	Low	Creamish	„	„	„
ENDB14	„	Moderate	Raised	No	Creamish	„	„	„
ENDB15	„	Pin point	Convex	Low	White	„	„	„
ENDB16	„	Pin point	„	Low	Creamish	„	„	„
ENDB17	„	Pin point	„	Low	Creamish	„	„	„

Antimicrobial activity of endophytic bacterial isolates

A total of seventeen bacterial endophytes were isolated from *Aloe vera* plant, all the isolates were subjected for antimicrobial screening. A total of sixteen isolates have shown antimicrobial activity against tested bacteria, 94.11% of the endophytic bacteria were active and shows prominent antimicrobial activity. Among all the bacterial endophytes isolates ENDB12 showed maximum antimicrobial activity against *E. coli* and lowest was found in ENDB 1 whereas ENDB4, 5, 8, 14, 16 and 17 have no inhibition activity against *E. coli*. Similarly, isolate ENDB8 has the highest antimicrobial activity against *S. pyogenes*, lowest was exhibited by ENDB11

and ENDB 16 and 17 has shown no antimicrobial activity against *S. pyogenes* and except ENDB14 (no zone of inhibition) rest all the 16 bacterial endophytes have shown good antimicrobial activity against acne bacterial isolate (ABI). ENDB3 showed maximum activity against ABI and lowest was displayed by ENDB13. Although, the bacterial endophytes have shown promising results against fungal pathogens, isolate ENDB10 has shown highest activity against *A. niger*, lowest was exhibited by ENDB5 and ENDB14,15,16 and 17 have exhibited no activity. Similarly, ENDB10 showed maximum antimicrobial activity against *Fusarium oxysporum*, ENDB 14, 15, 16, and 17 have displayed no activity whereas the ENDB12 shown lowest activity against *Fusarium oxysporum*. Therefore, it could be concluded that among all the tested microorganisms only, 64.7% (11/17) bacterial endophytes were found to be active against *E. coli*, Similarly, 88.23% (15/17), 94.11% (16/17), 76.4% (13/17) and 76.47% (13/17) bacterial endophytes were shown to be active against *S. pyogenes*, acne bacterial isolate (ABI), *A. niger* and *Fusarium oxysporum* respectively (Fig. 1, 2, 3, 4, and 5). Statistical significance ($p \leq 0.05$) of the data was obtained by Analysis of variance (One way ANOVA) for differences in means.




Figure 1. Antimicrobial potentiality spectra of bacterial endophytes against *E. coli*.




Figure 2. Antimicrobial potentiality spectra of bacterial endophytes against *S. pyogenes*.




Figure 3. Antimicrobial potentiality spectra of bacterial endophytes against ABI.




Figure 4. Antimicrobial potentiality spectra of bacterial endophytes against A. niger.




Figure 5. Antimicrobial potentiality spectra of bacterial endophytes against Fusarium oxysporum.

In the present investigation results of antibiotic susceptibility of obtained bacterial endophytes against standard penicillin G P(10 units), Oxacillin (1 mcg), Cephalathin (30 mcg), Clindamycin (2 mcg), Erythromycin (15 mcg) and Amoxyclav (30 mcg) were evaluated and the tested antibiotics displayed no inhibitory effect on the growth of bacterial endophytes therefore it could be concluded that at the given concentration all the bacterial endophytes were resistant to all the used antibiotics (see Table 2).

Table 2. Antibiogram of bacterial endophytes against six standard antibiotics.

Bacterial Endophyte	Penicillin G P (10 units)	Oxacillin (1 mcg)	Cephalathin (30 mcg)	Clindamycin (2 mcg)	Erythromycin (15 mcg)	Amoxyclav (30 mcg)
ENDB 1	-	-	-	-	-	-
ENDB 2	-	-	-	-	-	-
ENDB 3	-	-	-	-	-	-
ENDB 4	-	-	-	-	-	-
ENDB 5	-	-	-	-	-	-
ENDB 6	-	-	-	-	-	-
ENDB 7	-	-	-	-	-	-
ENDB 8	-	-	-	-	-	-
ENDB 9	-	-	-	-	-	-
ENDB 10	-	-	-	-	-	-
ENDB 11	-	-	-	-	-	-
ENDB 12	-	-	-	-	-	-
ENDB 13	-	-	-	-	-	-
ENDB 14	-	-	-	-	-	-
ENDB 15	-	-	-	-	-	-
ENDB 16	-	-	-	-	-	-
ENDB17	-	-	-	-	-	-

- = Resistant strains against used antibiotics.

3.1. Application in paper and textile dyeing

In the present investigation ENDB3 produced an extracellular brilliant green-brown colour pigment under submerged condition (Fig. 6). The obtained pigment showed good results in dyeing of paper, cotton and cotton fabric without using mordant, visibly mark difference observed after 24 hr of dyeing with extracted pigment (Fig. 7).





Figure 6. Pigment produced under submerged condition.



(a)



(b)



Figure 7. Dyed objects: (a) Paper: Pre (L) and After dyeing (R) (b) cotton: Pre (R) and After dyeing (L) and (c) Cloth: Pre (R) and After dyeing (L).

4. Discussion

Recent attraction has been focused to study the endophytic microbial flora of medicinal plants because they have wide range of applications and found as symbiont with the host plant, various have been studied the endophytic microbial flora of many medicinal plants. It is found that there is a huge diversity of endophyte microbes found in various parts of plant tissues. Similarly, the findings suggested by Strobel et al. [16] *Muscodora albus* a fungus from cinnamon tree and Lilley et al. [17] reported 114 endophytic isolates of sugar beet of 23 different genera, supported the presence of endophytes in plant tissues. In the present study we have obtained 17 different strains of bacterial endophytes from medicinal plant Aloe vera (see Table 1) and their antimicrobial potential was evaluated (Fig. 1, 2, 3, 4, and 5). Similar, studies have been reported by Akinsanya et al. [18] they studied the endophytic bacterial flora of Aloe vera plant and reported 29 different species from the plant simultaneously they also studied the antimicrobial spectra of the isolates and focused on the antimicrobial activities of bioactive molecules of bacterial endophytes of Aloe vera plant. Santos et al. [19] also studied the fungal endophytic flora of Aloe vera plant and their antimicrobial potentials against pathogens were reported. Endophytic microorganisms have many important properties because they are connected to their metabolic capabilities and can provide protection to the plant by producing variety of bioactive molecules [20, 21, and 22].

Assessment of antimicrobial potentials of bacterial endophytes of Aloe vera plant has been performed against bacterial and fungal pathogens. There seems to be antimicrobial effects on the growth of pathogens in terms of inhibition were tested (Fig. 1,2,3,4 and 5). Similarly, Huang et al. [23] Li and Strobel [24], and Sinha et al. [25] also studied the antimicrobial activities of plant endophytes. In this work isolated, total 17 bacterial endophytes from Aloe vera plant and their antimicrobial potential was assessed out of these 16 bacterial endophytes (94.11% of isolated bacterial endophytes) were shown to be active against *E. coli*, *S. pyogenes*, acne bacterial isolate, *A. niger* and *F. oxysporum* and inhibited the growth of tested microorganisms. Various have been suggested the similar result for example, 27.6% isolated microbial strains of *Camptotheca acuminata* showed antimicrobial activity against pathogens [26], similarly, 8.3% isolated microbial strains from *Aquilaria sinensis* and *Dracaena cambodiana* displayed antimicrobial activity against pathogens [27] whereas Akinsanya et al. [18] reported that endophytic bacterial flora of Aloe vera plant displayed about 48.3% antimicrobial activity against human pathogens and Jalgaonwala [28] suggested <10% antimicrobial activity was exhibited by bacterial endophytes against pathogenic microbes. Endophytic microorganisms ever been the largest source for bioactive secondary metabolites, for example, the endophytic strains of fungal genus *Nigrospora* reported as a source of bioactive molecules [29, 30, and 31]. In this work the best antifungal activity was found in ENDB10 against *F. oxysporum* and best antibacterial activity was displayed by ENDB3 against acne bacterial isolate. The present study suggested that the bacterial endophyte ENDB10 has shown good antifungal activity against both tested pathogenic fungi (*A. niger* and *F. oxysporum*) and ENDB14, 15, 16, and 17 have shown no activity against tested fungi (*A. niger* and *F. oxysporum*) whereas the bacterial endophyte ENDB3 has shown the maximum and wide spectrum of antimicrobial activity against the bacterial pathogen used.

Antibiogram of isolated bacterial endophytes against six standard antibiotics exhibited resistance against specific concentration of all the antibiotics (Penicillin, Oxacillin, Cephalathin, Clindamycin, Erythromycin and Amoxyclav) used (see Table 2). Burman et al. [32] reported the similar findings where bacterial isolates shown

resistant to standard antibiotics like ampicillin (10 µg/disc), penicillin G (10 µg/disc) and amoxicillin (30 µg/disc).

Natural pigments are highly recommended to overcome the adverse effects of artificial and synthetic colorants [33, 34] used worldwide. The synthetic colorants have toxic impacts to human as well as many adverse impacts to the environment [35]. Production of bacterial pigment is an emerging field with potential in different industrial sectors. Extracellular pigments are easy, cheap to obtain, ecofriendly and has potential to be used in textile dyeing [36]. In this study, the extracellular pigment was produced by bacterial endophyte ENDB3 under static condition since little studies of pigment production from bacterial endophytes have been reported till yet. Khanam and Chandra [37] also reported dye yielding endophytic bacteria from plant *Beta Vulgaris* L. Many authors have suggested that bacterial pigment and their exploration of dyeing potential in different industries [13, 38, and 39], they have also reported bacterial pigment production and their application in textile and other dyeing industries. The extracted pigment can be potentially used in various sectors of industries like textile [40], paper dyeing [40, 41, and 42], soap coloring, candles [42], food colorant [42] etc. In this work the bacterial pigment is used to dye fabric, cotton and papers, although Ahmad [42] also reported the similar use of bacterial pigment for dyeing fabric, soaps, candles and papers. The finding contributes to increase the industrial and pharmacological potential value of bioactive molecules of bacterial endophytes from Aloe vera plant.

Furthermore, observations indicates that the bacterial endophytes of Aloe vera plant have industrial potential as they produce bioactive molecules that could use as antimicrobial agent and natural colorant.

It could be concluded that the endophytic microbes of Aloe vera plant have medicinal and other industrial potentials. Hence, the study recommends further investigation is required of these bioactive molecules of bacterial endophytes from Aloe vera for dyeing and pharmacological industries.

5. Conclusion

The present investigation revealed that the bacterial endophytes of Aloe vera plant have wide variety of bioactive molecules that inhibit the growth of pathogenic micro-organism both bacteria and fungi. ENDB3 displayed best antibacterial activity against acne bacterial isolate whereas ENDB10 showed maximum antifungal activity against *F. oxysporum*. Bacterial endophytes reflected resistant antibiogram towards different standard antibiotics (at specific concentration) used. The isolate ENDB3 has shown very good antibacterial activity against acne bacterial isolate and the strain has also shown promising results in paper and textile dyeing. Therefore, we found that bacterial endophytes of Aloe vera have great industrial potential to be explored especially, in pharmacological and dyeing industries. This study has established bacterial endophyte ENDB3 as a cost-effective source of antimicrobial bioactive molecules and bacterial pigment that may have diverse industrial applications (in Pharma and Dyeing industries). Further studies are recommended to identify the bioactive molecules produced by ENDB3 and ENDB10 in order to discover new drugs with antibacterial, antifungal activity respectively. Hence, optimization studies are needed to explore the potential of ENDB3 in textile and paper dyeing effectively, large scale of production investigation of the bioactive molecules is recommended to be used as novel antibiotic and dyeing pigment which can open new opportunities for commercialization of the antibiotic and dyeing pigment bioactive molecules.

Acknowledgement

The authors are thankful to the director and department of Life Science, Cancer Hospital and Research Center for providing facilities to conduct the research.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] T. Reynolds and A. C. Dweck, "Aloe vera leaf gel: a review update," *J. Ethnopharmacol.*, vol. 68, no. 1, pp. 3–37, 1999, doi: [https://doi.org/10.1016/S0378-8741\(99\)00085-9](https://doi.org/10.1016/S0378-8741(99)00085-9).
- [2] M. D. Boudreau and F. A. Beland, "An evaluation of the biological and toxicological properties of *Aloe barbadensis* (Miller), *Aloe vera*," *J. Environ. Sci. Heal. - Part C Environ. Carcinog. Ecotoxicol. Rev.*, vol. 24, no. 1, pp. 103–154, 2006, doi: 10.1080/10590500600614303.

- [3] A. Snijders, "Medicinal plants of South Africa, 2nd edition," J. S. Afr. Vet. Assoc., vol. 81, no. 3, pp. 188–188, 2010.
- [4] N. C.A., A. L.A., and P. J.D., "Herbal medicines. A guide for health-care professionals." 1996.
- [5] T. L. Du, F. H. Van Der Westhuizen, and L. Botes, "Aloe ferox leaf gel phytochemical content, antioxidant capacity, and possible health benefits," J. Agric. Food Chem., vol. 55, no. 17, pp. 6891–6896, 2007, doi: 10.1021/jf071110t.
- [6] C. Costelloe, C. Metcalfe, A. Lovering, D. Mant, and A. D. Hay, "Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: Systematic review and meta-analysis," BMJ, vol. 340, no. 7756, p. 1120, 2010, doi: 10.1136/bmj.c2096.
- [7] G. Strobel and B. Daisy, "Bioprospecting for microbial endophytes and their natural products," Microbiol. Mol. Biol. Rev., vol. 67, no. 4, pp. 491–502, Dec. 2003, doi: 10.1128/MMBR.67.4.491-502.2003.
- [8] R. J. Rodriguez, J. F. White, A. E. Arnold, and R. S. Redman, "Fungal endophytes: Diversity and functional roles: Tansley review," New Phytol., vol. 182, no. 2, pp. 314–330, 2009, doi: 10.1111/j.1469-8137.2009.02773.x.
- [9] M. Caroch, P. Morales, and I. C. F. R. Ferreira, "Natural food additives: Quo vadis? - 1-s2.0-S0924224415001508-main.pdf," Nat. food Addit. Quo vadis?, vol. 1, p. 45, 2015, [Online]. Available: http://ac.els-cdn.com/S0924224415001508/1-s2.0-S0924224415001508-main.pdf?_tid=45d4048e-7655-11e5-9546-00000aab0f26&acdnat=1445254612_c9894fbf6530dd3d5290c37a51efea74.
- [10] M. Shahid, Shahid-Ul-Islam, and F. Mohammad, "Recent advancements in natural dye applications: A review," J. Clean. Prod., vol. 53, pp. 310–331, 2013, doi: 10.1016/j.jclepro.2013.03.031.
- [11] R. H. Kostick, "(12) Patent Application Publication (10) Pub . No . : US 2006 / 0280762 A1," vol. 1, no. 19, 2006.
- [12] E. Fernández-García, I. Carvajal-Lérida, M. Jarén-Galán, J. Garrido-Fernández, A. Pérez-Gálvez, and D. Hornero-Méndez, "Carotenoids bioavailability from foods: From plant pigments to efficient biological activities," Food Res. Int., vol. 46, no. 2, pp. 438–450, 2012, doi: 10.1016/j.foodres.2011.06.007.
- [13] C. K. Venil, Z. A. Zakaria, and W. A. Ahmad, "Bacterial pigments and their applications," Process Biochem., vol. 48, no. 7, pp. 1065–1079, 2013, doi: 10.1016/j.procbio.2013.06.006.
- [14] A. W. Bauer, W. M. Kirby, J. C. Sherris, and M. Turck, "Antibiotic susceptibility testing by a standardized single disk method.," Tech. Bull. Regist. Med. Technol., vol. 36, no. 3, pp. 49–52, 1966, doi: 10.1007/978-94-009-0753-9_4.
- [15] M. M. Cowan, "Plant products as antimicrobial agents," Clin. Microbiol. Rev., vol. 12, no. 4, pp. 564–582, Oct. 1999, doi: 10.1128/CMR.12.4.564.
- [16] G. A. Strobel, "Endophytes as sources of bioactive products," Microbes Infect., vol. 5, no. 6, pp. 535–544, 2003, doi: 10.1016/S1286-4579(03)00073-X.
- [17] A. K. Lilley, J. C. Fry, M. J. Bailey, and M. J. Day, "Comparison of aerobic heterotrophic taxa isolated from four root domains of mature sugar beet (*Beta vulgaris*)," FEMS Microbiol. Ecol., vol. 21, no. 3, pp. 231–242, 2006, doi: 10.1111/j.1574-6941.1996.tb00350.x.
- [18] M. A. Akinsanya, J. K. Goh, S. P. Lim, and A. S. Y. Ting, "Diversity, antimicrobial and antioxidant activities of culturable bacterial endophyte communities in Aloe vera," FEMS Microbiol. Lett., vol. 362, no. 23, 2015, doi: 10.1093/femsle/fnv184.
- [19] I. P. dos Santos, L. C. N. da Silva, M. V. da Silva, J. M. de Araújo, M. da S. Cavalcanti, and V. L. de M. Lima, "Antibacterial activity of endophytic fungi from leaves of *Indigofera suffruticosa* Miller (Fabaceae)," Front. Microbiol., vol. 6, no. MAY, pp. 1–7, 2015, doi: 10.3389/fmicb.2015.00350.
- [20] B. J. Duijff, V. Gianinazzi-Pearson, and P. Lemanceau, "Involvement of the outer membrane lipopolysaccharides in the endophytic colonization of tomato roots by biocontrol *Pseudomonas fluorescens* strain WCS417r," New Phytol., vol. 135, no. 2, pp. 325–334, 1997, doi: 10.1046/j.1469-8137.1997.00646.x.
- [21] R. X. Tan and W. X. Zou, "Endophytes: A rich source of functional metabolites," Nat. Prod. Rep., vol. 18, no. 4, pp. 448–459, 2001, doi: 10.1039/b100918o.
- [22] X. Lin, C. Lu, Y. Huang, Z. Zheng, W. Su, and Y. Shen, "Endophytic fungi from a pharmaceutical plant, *Camptotheca acuminata*: Isolation, identification and bioactivity," World J. Microbiol. Biotechnol., vol. 23, no. 7, pp. 1037–1040, 2007, doi: 10.1007/s11274-006-9329-8.

- [23] J. Y. Li and G. A. Strobel, "Jesterone and hydroxy-jesterone antioomycete cyclohexenone epoxides from the endophytic fungus *Pestalotiopsis jesteri*," *Phytochemistry*, vol. 57, no. 2, pp. 261–265, 2001, doi: 10.1016/S0031-9422(01)00021-8.
- [24] A. Sinha, R. Priya, M. Nimisha, and W. Jabez Osborne, "Impact of endophytic *Ralstonia* sp. From *Aloe vera* gel and its antimicrobial activity," *Asian J. Pharm. Clin. Res.*, vol. 8, no. 1, pp. 259–262, 2015.
- [25] X. Lin, C. Lu, Y. Huang, Z. Zheng, W. Su, and Y. Shen, "Endophytic fungi from a pharmaceutical plant, *Camptotheca acuminata*: Isolation, identification and bioactivity," *World J. Microbiol. Biotechnol.*, vol. 23, no. 7, pp. 1037–1040, 2007, doi: 10.1007/s11274-006-9329-8.
- [26] L. J. Gong and S. X. Guo, "Endophytic fungi from *Dracaena cambodiana* and *Aquilaria sinensis* and their antimicrobial activity," *African J. Biotechnol.*, vol. 8, no. 5, pp. 731–736, 2009, doi: 10.4314/ajb.v8i5.59937.
- [27] R. E. Jalgaonwala, B. V. Mohite, and R. T. Mahajan, "Evaluation of Endophytes for their Antimicrobial activity from Indigenous Medicinal Plants belonging to North Maharashtra region India," *Int. J. Pharm. Biomed. Res.*, vol. 1, no. 5, pp. 136–141, 2010.
- [28] T. Fukushima, M. Tanaka, M. Gohbara, and T. Fujimori, "Phytotoxicity of three lactones from *Nigrospora sacchari*," *Phytochemistry*, vol. 48, no. 4, pp. 625–630, 1998, doi: 10.1016/S0031-9422(97)01023-6.
- [29] J. C. Kim, G. J. Choi, J. H. Park, H. T. Kim, and K. Y. Cho, "Activity against plant pathogenic fungi of phomalactone isolated from *Nigrospora sphaerica*," *Pest Manag. Sci.*, vol. 57, no. 6, pp. 554–559, 2001, doi: 10.1002/ps.318.
- [30] J. H. Zhao, Y. L. Zhang, L. W. Wang, J. Y. Wang, and C. L. Zhang, "Bioactive secondary metabolites from *Nigrospora* sp. LLGLM003, an endophytic fungus of the medicinal plant *Moringa oleifera* Lam," *World J. Microbiol. Biotechnol.*, vol. 28, no. 5, pp. 2107–2112, 2012, doi: 10.1007/s11274-012-1015-4.
- [31] S. Burman, K. Bhattacharya, D. Mukherjee, and G. Chandra, "Antibacterial efficacy of leaf extracts of *Combretum album* Pers. against some pathogenic bacteria," *BMC Complement. Altern. Med.*, vol. 18, no. 1, pp. 1–8, 2018, doi: 10.1186/s12906-018-2271-0.
- [32] D. Manikprabhu and K. Lingappa, "γ Actinorhodin a natural and attorney source for synthetic dye to detect acid production of fungi," *Saudi J. Biol. Sci.*, vol. 20, no. 2, pp. 163–168, 2013, doi: 10.1016/j.sjbs.2013.01.004.
- [33] B. Padhan, K. Poddar, D. Sarkar, and A. Sarkar, "Production, purification, and process optimization of intracellular pigment from novel psychrotolerant *Paenibacillus* sp. BPW19," *Biotechnol. Reports*, vol. 29, p. e00592, 2021, doi: 10.1016/j.btre.2021.e00592.
- [34] S. Babitha, "Microbial pigments," *Biotechnol. Agro-Industrial Residues Util. Util. Agro-Residues*, pp. 147–162, 2009, doi: 10.1007/978-1-4020-9942-7_8.
- [35] Z. Chadni, M. H. Rahaman, I. Jerin, K. M. F. Hoque, and M. A. Reza, "Extraction and optimisation of red pigment production as secondary metabolites from *Talaromyces verruculosus* and its potential use in textile industries," *Mycology*, vol. 8, no. 1, pp. 48–57, 2017, doi: 10.1080/21501203.2017.1302013.
- [36] B. Khanam and R. Chandra, "Isolation and identification of endophytic bacteria producing bright red pigment from the dye yielding plant *beta vulgaris* L," *Int. J. Pharm. Pharm. Sci.*, vol. 7, no. 5, pp. 220–224, 2015.
- [37] A. Shirata et al., "Isolation of bacteria producing bluish-purple pigment and use for dyeing," *Japan Agric. Res. Q.*, vol. 34, no. 2, pp. 131–140, 2000.
- [38] R. Poorniammal, M. Parthiban, S. Gunasekaran, R. Murugesan, and G. Thilagavathi, "Natural dye production from *Thermomyces* sp fungi for textile application," *Indian J. Fibre Text. Res.*, vol. 38, no. 3, pp. 276–279, 2013.
- [39] S. A. S. Mapari, K. F. Nielsen, T. O. Larsen, J. C. Frisvad, A. S. Meyer, and U. Thrane, "Exploring fungal biodiversity for the production of water-soluble pigments as potential natural food colorants," *Curr. Opin. Biotechnol.*, vol. 16, no. 2, pp. 231–238, 2005, doi: 10.1016/j.copbio.2005.03.004.
- [40] R. Bott, "Brock Biology of Microorganisms, 14th Edition- Madigan," *Igarss* 2014, no. 1, pp. 1–5, 2014.
- [41] W. A. Ahmad, W. Y. W. Ahmad, Z. A. Zakaria, and N. Z. Yusof, "Application of Bacterial Pigments as Colorant," pp. 57–74, 2012, doi: 10.1007/978-3-642-24520-6_4.
- [42] K. Malik, J. Tokkas, and S. Goyal, "Microbial Pigments: A review," no. May, pp. 361–365, 2012.